

Fabian A. Ramos, M.D.
John J. D'Auria, M.D.
José Sarria, M.D.

100 3rd Ave. West, Suite #110 Bradenton, FL 34205
5741 Bee Ridge Road Suite 550 Sarasota, FL 34233
Phone: (941) 708-9555 Fax: (941) 708-5465



MAC Anesthesia or Light Sedation – Post Op Care

Special Instructions AFTER your procedure

- Patients who received sedation or anesthesia should return home from the office **immediately upon discharge** and lie down (bed/recliner) until fully recovered from any side effects of the anesthetic. Anesthetic effects vary by the individual and you may feel drowsy for a short period of time and rarely for several hours (we use, in most cases, a very short-lived sedative/anesthetic).
- Do not use appliances or equipment that could be dangerous, such as power tools, stoves, burners, lawnmowers, and/or garbage disposals, etc.
- Be aware of dizziness. Take your time and walk slowly as sudden changes of position may also cause nausea.
- You should not operate any mechanical equipment or drive a motor vehicle for at least 24 hours or longer.
- You should not make important decisions (as you may change your mind the following day).
- Do not drink any alcoholic beverages the day of the procedure as the anesthetic causes an exaggerated reaction to the alcohol which could be very dangerous.
- **DIET:** For the most part, you may gradually return to a regular diet two hours after the procedure (use common sense). If you feel nauseated or sick to your stomach, you should drink small sips of clear liquid (broth, apple juice, ginger ale, tea or Jell-O). If these liquids do not make you sick to your stomach, try eating soft foods like potatoes, rice and pasta. Eat any nourishing foods that can be eaten with comfort. After procedures, diabetics need to maintain their normal diet as much as possible and follow your physician's instructions regarding your insulin schedule. Proper nourishment aids in the healing process.
- **HYGIENE (applies only for needle procedures. If you have a scar/sutures you will have different instructions):** Since you have showered prior to the procedure, please do not disturb the surgical area today. Leave the dressings or bandages as placed unless they are soaked with drainage. For the most part you should not need to replace these bandages, unless some oozing continues (applies only to limited types of procedures). The next day you may shower with soap and water and pat the area dry, no rubbing. **DO NOT** submerge your body in water (bath, pool or spa) for 48 hours following the needle procedure.
- These are general guidelines and some procedures have specific post-operative recovery time. As a general rule, some pain (minor) may present within 24 hours after the procedure in the injection site and may last up to 48 hours. Use ice and rest techniques appropriately.
- **Please call our office at extension 120**, if there are any unexpected symptoms. Use common sense and If necessary you may go to the emergency room for unexpected severe pain or weakness. If the needle or surgical site becomes red, inflamed, draining occurs or if you become febrile, please come to our office for examination.